Carbohydrate Research, 255 (1994) C5–C6
Elsevier Science B.V., Amsterdam

CONTENTS

Foreword vii

The effect of variation of substitution on the solution conformation of heparin: a spectroscopic and molecular modelling study
B. Mulloy, M.J. Forster, C. Jones (Potters Bar, United Kingdom), A.F. Drake (London, United Kingdom), E.A. Johnson (Potters Bar, United Kingdom) and D.B. Davies (London, United Kingdom) 1

A reversed phase HPLC method for the simultaneous determination of all monosaccharides contained in galactosaminoglycan isomers from human aorta proteoglycans
M. Formato, A. Senes, F. Soccolini, R. Coinu and G.M. Cherchi (Sassari, Italy) 27

Preliminary investigation into the purification, NMR analysis, and molecular modelling of chondroitin sulphate epitopes
S.M.T. D'Arcy, S.L. Carney and T.J. Howe (Windlesham, United Kingdom) 41

Synthesis of hyaluronic acid-related di-, tri-, and tetra-saccharides having an *N*-acetylglucosamine residue at the reducing end
T.M. Slaghek (Utrecht, Netherlands), Y. Nakahara, T. Ogawa (Saitama, Tokyo, Japan), J.P. Kamerling and J.F.G. Vliegenthart (Utrecht, Netherlands) 61

Substrate specificities of glycosyltransferases involved in formation of heparin precursor and *E. coli* K5 capsular polysaccharides
K. Lidholt, M. Fjelstad (Uppsala, Sweden), K. Jann (Freiburg, Germany) and U. Lindahl (Uppsala, Sweden) 87

Synthesis of sulfated and phosphorylated glycopeptides from the carbohydrate–protein linkage region of proteoglycans
S. Rio, J.-M. Beau and J.-C. Jacquinet (Orléans, France) 103

Preliminary chemical, biochemical, and pharmacological characterization of a low molecular weight dermatan sulphate
G.P. Ferrari, D. Marchesini and A.P. Maggi (Milan, Italy) 125

Dermatan sulfate from beef mucosa: structure, physicochemical and biological properties of fractions prepared by chemical depolymerization and anion-exchange chromatography
N. Volpi (Modena, Italy) 133

Structural studies on the chondroitinase ABC-resistant sulfated tetrasaccharides isolated from various chondroitin sulfate isomers
K. Sugahara, K. Shigeno, M. Masuda (Kobe, Japan), N. Fujii, A. Kurosaka (Kyoto, Japan) and K. Takeda (Kobe, Japan) 145

Chondroitinase ABC-resistant sulfated trisaccharides isolated from digests of chondroitin/dermatan sulfate chains
K. Sugahara, Y. Takemura, M. Sugiura (Kobe, Japan), Y. Kohno, K. Yoshida (Tokyo, Japan), K. Takeda (Kobe, Japan), K.H. Khoo, H.R. Morris and A. Dell (London, United Kingdom) 165

Evidence of a selective free radical degradation of heparin, mediated by cupric ion
Z. Liu and A.S. Perlin (Montreal, Canada). 183

Molecular weight of low molecular weight heparins by ^{13}C nuclear magnetic resonance spectroscopy
U.R. Desai and R.J. Linhardt (Iowa City, IA, USA) 193

Isolation and partial characterization of a novel amino sugar-containing fucan sulfate from commercial *Fucus vesiculosus* fucoidan
T. Nishino, C. Nishioka, H. Ura and T. Nagumo (Kitasato, Japan) 213

A sulfated α-L-fucan from sea cucumber
A.-C. Ribeiro, R.P. Vieira, P.A.S. Mourão (Rio de Janeiro, Brazil) and B. Mulloy (Potters Bar, United Kingdom) 225

The sulphation pattern in chondroitin sulphate chains investigated by chondroitinase ABC and ACII digestion and reactivity with monoclonal antibodies
T.E. Hardingham, A.J. Fosang, N.J. Hey, P.K. Hazell, W.J. Kee and R.J.F. Ewins (London, United Kingdom) 241

The relationship between proteoglycan synthesis in Swarm chondrocytes and pathways of cellular energy and UDP-sugar metabolism
R.M. Mason and C. Sweeney (London, United Kingdom) 255

Notes

Preparation, purification, and structural characterization of linear oligogalacturonides. An FAB-mass spectrometric and NMR spectroscopic study
V.-M. Ló, M.G. Hahn and H. Van Halbeek (Athens, GA, USA) 271

A procedure for the analysis of glycosaminoglycan mixtures based on digestion by specific enzymes
G.C. Slim, R.H. Furneaux (Lower Hutt, New Zealand) and S.C. Yorke (Palmerston North, New Zealand) 285

Synthesis of the basic disaccharide unit of heparin
G. Jaurand, C. Tabeur and M. Petitou (Gentilly, France) 295

600 MHz ^{1}H NMR study of a fucose-containing heptasaccharide derived from a keratanase digestion of bovine articular cartilage keratan sulphate
G.-H. Tai, T.N. Huckerby and I.A. Nieduszynski (Lancaster, United Kingdom) 303

Author index C1

Subject index C3